

**FANGTONG**  
**Caltong**

Calcium Borogluconate Injection

**Composition**

Calcium gluconate ..... 265mg  
Boric acid.....35mg  
Magnesium chlorid.....40mg

**Indications**

As a source of calcium for cattle, sheep and swine, to aid in the treatment of hypocalcemia, e.g., milk fever in cattle.

**Directions**

Administer by slowly intravenous or subcutaneous injection.  
Intravenous administration is recommended for the treatment of acute conditions.  
When administered subcutaneously, the dose should be divided over several sites.  
Warm the solution to body temperature prior to use.  
Use aseptic technique for administration.  
Dosage may be repeated after several hours if required or as recommended by a veterinarian.

**Dosage**

For one dose, cattle, sheep and swine: 10ml per 24kg body weight or as recommended by a veterinarian.

**Cautions**

Use only if solution is clear.  
If no improvement is noticed in the condition being treated after 24 hours or less depending on the severity of the condition, the diagnosis should be re-evaluated and a veterinarian consulted.  
Do not retain partially used vial.

**Storage**

Do not store above 20°C, protect from light.

**Shelf Life**

3 years.



**For Veterinary Use Only**  
**Keep Out of Reach of Children**



Tech consultation Tech on facebook  
EDITION NO.: CAL 202505 VERIFY DATE: since May, 2025

100\*130mm