

 **FANGTONG**

Doratong 3.15%

Doramectin Injection

Composition

Each ml contains Doramectin 31.5 mg.

Indications

Broad spectrum endectocide. For the control and treatment of endo and ectoparasites in bovines. This product has a slow release and absorption mechanism that provides a prolonged efficacy, eliminating the parasites and preventing new infestations for long periods, achieving an effective treatment with prophylactic effect.
Good overall efficiency (90 – 95%) until the day 60 post treatment against gastrointestinal parasites.

Dosage and Administration

Administer by subcutaneous route and given a dose of 630 µg/kg body weight. Each 1ml/ 50 kg of body weight, in front or behind the scapula, taking into account the aseptic measures that corresponding to the application of injectable.
Do not administer more than 10 ml per inoculation site.
In fields where is planned the tick eradication, administer every 45 days.

Withdrawal Period

Meat: 120 days. Do not administer in lactating dairy cows when the milk is intended for human consumption.

Storage

Do not store above 30°C, protect from light.

Validity

3 years.

Presentation

Available in 50ml or 100ml bottle.

Manufactured by

Chongqing Fangtong Animal Pharmaceutical Co., Ltd
No. 80, East Part of Changzhou Road, Rongchang, Chongqing, China

MOCCA REG NO.: AUD-APH-10-2918431



For Veterinary Use Only
Keep Out of Reach of Children



EDITION NO.: 2409 DORA VERIFY DATE: since Sep, 2024

100*130mm, 5500张