

Category: POM

Ivertong Super



INJECTABLE SOLUTION OF
IVERMECTIN 1%w/v, CLORSULON 10%w/v

COMPOSITION

Each ml contains ivermectin 10 mg, clorsulon 100 mg.

INDICATIONS

Ivertong Super is indicated for the effective treatment and control of the following parasites of cattle:

Gastrointestinal Roundworms (adults and fourth-stage larvae):

Ostertagia ostertagi (including inhibited *O. ostertagi*)

O. lyrata

Haemonchus placei

Trichostrongylus axei

T. colubriformis

Cooperia oncophora

C. punctata

C. pectinata

Bunostomum phlebotomum

Nematodirus helvetianus (adults only)

N. spathiger (adults only)

Oesophagostomum radiatum

Lungworms (adults and fourth-stage larvae):

Dictyocaulus viviparus

Liver Flukes:

Fasciola hepatica (adults only)

Cattle Grubs (parasitic stages):

Hypoderma bovis

H. lineatum

Sucking Lice:

Linognathus vituli

Haematopinus eurysternus

Solenopotes capillatus

Mange Mites (cattle scab*):

Psoroptes ovis (syn. *P. communis* var. *bovis*)

Sarcoptes scabiei var. *bovis*

Persistent Activity

Ivertong Super has been proved to effectively control infections and to protect cattle from reinfection with *Dictyocaulus viviparus* and *Oesophagostomum radiatum* for 28 days after treatment; *Ostertagia ostertagi*, *Trichostrongylus axei* and *Cooperia punctata* for 21 days after treatment; *Haemonchus placei*, and *Cooperia oncophora* for 14 days after treatment.

DOSAGE AND ADMINISTRATION

Ivertong Super should be administered only by subcutaneous injection at the recommended dosage level of 1 ml / 50 kg body weight (based on a dosage level of 200 mcg ivermectin plus 2 mg clorsulon per kg body weight) under the loose skin in front of, or behind the shoulder.

FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN

100*130mm

ADVERSE REACTIONS

Transitory discomfort has been observed in some cattle following subcutaneous administration. Soft tissue swellings may occur at the site of injection. These reactions resolve over time without treatment.

CONTRAINDICATIONS

This product has been developed specifically for use in cattle only. Do not use in other animal species as severe adverse reactions, including fatalities in dogs, may occur. Do not use intramuscularly or intravenously. Do not use in animals with known hypersensitivity to the active ingredient or any of the excipients.

PRECAUTIONS

Divide doses greater than 10 mL between two injection sites to reduce occasional discomfort or site reaction. Different injection sites should be used for other parenteral products.

WARNINGS

Free ivermectin may adversely affect fish and some waterborne organisms on which they feed.

Do not contaminate water by direct application or the improper disposal of drug containers. Dispose of containers in an approved landfill or by incineration.

WITHDRAWAL PERIOD

Do not treat cattle within 21 days of slaughter.

Because a withdrawal time in milk has not been established, do not use in cattle producing milk for human consumption, do not use in female dairy cattle of breeding age.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

PRESENTATION

Available in 50 ml and 100ml bottle.

STORAGE

Store in a cool and dry place below 30°C, protect from light.

VALIDITY

3 years.

Manufactured by

FANGTONG

Chongqing Fangtong Animal Pharmaceutical Co.,Ltd

No. 80, East Part of Changzhou Road, Rongchang, Chongqing, China



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN

EDITION NO.: 2012TYL0XY VERIFY DATE: Since Dec., 2020

100*130mm