

Category: POM

FANGTONG

Nitrotong 34

Nitroxinil Injection 34%

Composition
Each ml contains Nitroxinil 340mg.

Pharmacology
Nitrotong 34 is nitrate derivative from benzene compounds and presents a flukicide effect and is commonly used and the one with best spectrum from all the nitrate derivative trematocides. The mechanism of action corresponds to a respiratory metabolism interference of the helminth, blocking the energy production by mitochondrial oxidative phosphorylation. Nitrotong 34 impedes to use the oxide-reduction reactions for adenosine triphosphate (ATP) production. After the use of all its energetic reserves the parasites die due to starvation.
Nitrotong 34 also causes the scolex detachment of the intestinal mucosa and the proximal proglottid of the cestode.

Indication
Nitrotong 34 is indicated for control of Fasciolosis (larval and adult stage) caused by Fasciola hepatica and F.gigantica, gastrointestinal parasitism caused by Haemonchus spp., Bunostomum spp., Oesophagostomum spp., Oestrus spp. in cattle, sheep, goats and camel, as well as Ancylostoma spp. and Uncinaria spp. in dogs.

Dosage and Administration
By subcutaneous injection only.
Normal aseptic precautions should be observed.
The Standard dosage is 10mg nitroxinil per kg bodyweight (=1.5 ml of Nitrotong /50kg b.w.).
If animals are to be treated collectively rather than individually, they should be grouped according to their body weight and dosed accordingly, in order to avoid under or overdosing.

Contraindications
Do not use in animals with known hypersensitivity to the active ingredient.
Do not exceed stated dose.

Over dosage
In the event of accidental over dosage, the symptoms are pyrexia, rapid respiration and increased excitability. Patients should be kept cool, and dextrose saline should be administered intravenously.

Adverse Reactions
Small swellings are occasionally observed at the injection site in cattle. These can be avoided by injecting the dose in two separate sites and massaging well to disperse the solution.
No systemic ill effects are to be expected when animals are treated at normal dosage.

Precautions
-Only dose cows on dry period.
-Do not mix in the same syringe or container with any other substance different to the product. The containers and any residue of product should be eliminated in a safe way(burial or incineration).
-The use on other species than authorized is not recommended.
-It can be used at any stage of pregnancy (however, in the last third, the product must be managed very carefully under professional supervision); it does not affect fertility, gestation nor fetus formation, and it does not affect the reproductive performance of stallions.
-In case of intoxication, keep the patient in a cold place and administer dextrose by endovenous route.
-In case of ingestion, wash mouth and request help to the veterinarian.
-Be careful of auto inject accidentally. Seek medical attention in case of accidental injection.

Withdrawal Period
Meat: 30 days.
Milk: 5 days.

Presentation
In 100ml or 250ml bottle.

Storage Condition
Store in a cool and dry place not above 30°C, protect from light.
Keep out of reach of children.

Shelf Life
2 years.

Manufacturer
Chongqing Fangtong Animal Pharmaceutical Co., Ltd
No. 80, East Part of Changzhou Road, Rongchang, Chongqing, China

Exclusive Agent in Iraq
Bilad Al-Rajih Company
For General Trading L.L.C

FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN



EDITION NO.: 2407NTR VERIFY DATE: since JULY, 2024

100*130mm 2000张