

Category: POM

Parvatong

Parvaquone Injection 15%



Composition

Each ml contains parvaquone 150mg.

Character

A clear ruby coloured, slightly viscous solution.

Indications

Treatment of East Coast fever (ECF, Theileria parva infection) of cattle. Parvaquone kills the schizont and piroplasm stages of parasite. The product may be used in animal showing clinical signs of theileriosis and infected animals which have not yet displayed clinical signs.

Dosage and administration

Injected by the intramuscular route into the neck muscles.

Calves: 1ml/15kg B.W.T. (10mg parvaquone per kg)

Adults:	Bodyweight	Dose
	100kg	7ml
	150kg	10ml
	200kg	14ml
	250kg	17ml
	300kg	20ml

Side effect

Local swelling may occur at the injection site but it is generally of a transient nature.

Warnings and precautions

1. In case of exceptionally severe infection, further treatment at the recommended dosage rate may be required at 48h intervals.
2. Concurrent infections such as anaplasmosis and pneumonia should be treated specially.
3. Do not use by the intravenous or subcutaneous route.

Withdrawal period

Meat - 28 days

Milk - 14 days

Presentation

Available in 50ml and 100ml bottle.

Storage conditions

Do not store above 30°C, store in a dry place. Protect from light. Keep out of reach of children.

Shelf life

3 years.

Use within 28 days after first opening.

TMDA REG. NO.:

Manufactured by

FANGTONG

Chongqing Fangtong Animal Pharmaceutical Co., Ltd
Banqiao Industrial Park, Rongchang, Chongqing, China

Distributed by: Anicrop Services Ltd



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN

EDITION NO. :2412PARVA VERIFY DATE: since DEC., 2024

Leaflet Size: 100×130 mm