

Category: POM

Penstong Duo LA

FANGTONG

Injectable Suspension of

Penicillin G procaine 100,000 IU/ml, Penicillin G benzathine 100,000 IU/ml
Dihydrostreptomycin sulphate 200 mg/ml

Composition

Each ml contains:
Penicillin G procaine 100,000 IU
Penicillin G benzathine 100,000 IU
Dihydrostreptomycin sulphate 200 mg

Description

Penicillin G procaine's fast action is enhanced with the long-acting properties of penicillin G benzathine, which is active during 5 days. The spectrum of activity is widened by the dihydrostreptomycin sulphate. Penicillin G procaine and penicillin G benzathine are small-spectrum penicillins with a bactericidal action against mainly Gram-positive bacteria like Clostridium, Corynebacterium, Erysipelothrix, Listeria, penicillinase-negative Staphylococcus and Streptococcus spp. Dihydrostreptomycin is an aminoglycoside with a bactericidal action against mainly Gram-negative bacteria like E. coli, Campylobacter, Klebsiella, Haemophilus, Pasteurella and Salmonella spp.

Indications

Arthritis, mastitis and gastrointestinal, respiratory and urinary tract infections caused by penicillin and dihydrostreptomycin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Klebsiella, Listeria, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp.

Contraindications

Hypersensitivity to penicillins, procaine and/or aminoglycosides.
Administration to animals with a seriously impaired renal function.
Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

Side effects

Side effects that may occur are ototoxicity, neurotoxicity, nephrotoxicity and hypersensitivity reactions.

Dosage and administration

For intramuscular administration only:
General dose: 1 ml per 10 kg body weight every 72 hours, or upon recommendation by a veterinarian every 48 hours.

Note: Shake well before use. Do not administer more than 20 ml in cattle and more than 5 ml in calves, sheep and goats per injection site.

Withdrawal periods

- For kidney : 45 days.
- For meat : 30 days.
- For milk : 5 days.

Presentation

Available in 50 ml and 100 ml bottle.

Storage

Store in a cool and dry place below 25°C, protect from light.
The product should be used immediately after opening.

Validity

2 years.

Manufactured by

Chongqing Fangtong Animal Pharmaceutical Co., Ltd
No. 80, East Part of Changzhou Road, Rongchang, Chongqing, China

MOCCA Registration No.: DXB-APH-10-2585352



FOR VETERINARY USE ONLY
KEEP OUT OF REACH OF CHILDREN
EDITION NO.:2308PEN Duo VERIFY DATE: since Aug, 2023

100*130mm, 6000张